



ELSEVIER

Contents lists available at ScienceDirect

## Human Movement Science

journal homepage: [www.elsevier.com/locate/humov](http://www.elsevier.com/locate/humov)



# Effects of whole body vibration on spasticity and lower extremity function in children with cerebral palsy



Hsin-Yi Kathy Cheng<sup>a,1</sup>, Yan-Ying Ju<sup>b,2</sup>, Chia-Ling Chen<sup>a,c,3</sup>,  
Li-Ling Chuang<sup>d,4</sup>, Chih-Hsiu Cheng<sup>d,\*</sup>

<sup>a</sup> Graduate Institute of Early Intervention, College of Medicine, Chang Gung University, Tao-Yuan, Taiwan

<sup>b</sup> Department of Adapted Physical Education, National Taiwan Sport University, Tao-Yuan, Taiwan

<sup>c</sup> Department of Physical Medicine and Rehabilitation, Chang Gung Memorial Hospital, Tao-Yuan, Taiwan

<sup>d</sup> Department and Graduate Institute of Physical Therapy, College of Medicine, Chang Gung University, Tao-Yuan, Taiwan

### ARTICLE INFO

#### Article history:

Available online 24 November 2014

#### PsycINFO classification:

3380

#### Keywords:

Cerebral palsy

Spasticity

Whole body vibration

Ambulation

Rehabilitation

### ABSTRACT

**Aim:** The aim of this study was to evaluate the effect of whole body vibration (WBV) on lower extremity spasticity and ambulatory function in children with cerebral palsy (CP) with a complete cross-over design.

**Method:** Sixteen participants aged 9.8(2.3) years received a 20-min WBV and a control condition in a counterbalanced order on two separate days. Change scores of each outcome variable were used to show the improvement.

**Results:** Repeated-measures analyses revealed significant differences in condition scores among variables including active range-of-motion (active ROM, increased), relaxation index (RI, increased), Modified Ashworth Scale (MAS, decreased), timed up-and-go (TUG, decreased), and Six Minute Walk Test (6MWT, increased). Significant differences were also found in time change scores for MAS

\* Corresponding author at: Department of Physical Therapy and Graduate Institute of Rehabilitation Science, College of Medicine, Chang Gung University, Tao-Yuan 333, Taiwan. Tel.: +886 3 211 8800x3714; fax: +886 3 211 8700.

E-mail addresses: [kcheng@mail.cgu.edu.tw](mailto:kcheng@mail.cgu.edu.tw) (H.-Y.K Cheng), [yanju@ntsuo.edu.tw](mailto:yanju@ntsuo.edu.tw) (Y.-Y. Ju), [clingchen@gmail.com](mailto:clingchen@gmail.com) (C.-L. Chen), [lchuang@mail.cgu.edu.tw](mailto:lchuang@mail.cgu.edu.tw) (L.-L. Chuang), [chcheng@mail.cgu.edu.tw](mailto:chcheng@mail.cgu.edu.tw) (C.-H. Cheng).

<sup>1</sup> Address: Graduate Institute of Early Intervention, College of Medicine, Chang Gung University, 259 Wen-Hua 1st Rd., Kwei-Shan, Tao-Yuan 333, Taiwan. Tel.: +886 3 211 8800x3667; fax: +886 3 211 8700.

<sup>2</sup> Address: Department of Adapted Physical Education, National Taiwan Sport University, 250 Wen-Hua 1st Rd., Kwei-Shan, Tao-Yuan 333, Taiwan.

<sup>3</sup> Address: Graduate Institute of Early Intervention, College of Medicine, Chang Gung University, Tao-Yuan 333, Taiwan.

<sup>4</sup> Address: Department of Physical Therapy and Graduate Institute of Rehabilitation Science, College of Medicine, Chang Gung University, Tao-Yuan 333, Taiwan.

<http://dx.doi.org/10.1016/j.humov.2014.11.003>

0167-9457/© 2014 Elsevier B.V. All rights reserved.

and 6MWT. Correlation results revealed that TUG was significantly correlated with RI ( $r = -.512$ ,  $p = .042$ ), and 6MWT ( $r = -.700$ ,  $p = .003$ ).

*Interpretation:* This study suggested that WBV intervention can control the spasticity, enhance ambulatory performance and increase active ROM. Along with previous results, data from this study revealed the potential use of WBV in clinical rehabilitation in children with CP. Future investigations should focus on finding the combination of treatment frequency and duration to achieve an ideal result.

© 2014 Elsevier B.V. All rights reserved.

---

## 1. Introduction

Cerebral palsy (CP), which occurs in 2–3 of every 1000 individuals, is one of the leading causes of movement and posture disorders (Koman, Smith, & Shilt, 2004). Spasticity is a frequent-observed phenomenon after upper motor neuron lesion, including stroke, head injury, spinal cord injury and CP, etc. About seventy to eighty percent of children with CP demonstrate spastic clinical features (Kriger, 2006). These children are affected by neuromuscular symptoms such as impaired sensation and increased muscle tone, and therefore demonstrate a decreased ability to voluntarily control their muscle activities. As a result, a wide range of movement dysfunction can occur.

The goals of spasticity management are to improve flexibility and movement ability. Clinically, other than the medication and injection treatments, muscle stretching and functional training have been considered an integral part of patient's daily management of spasticity. Our group has performed a series of studies investigating the benefits of repetitive passive movements onto spasticity control, sensory normalization and functional improvement (Cheng, Ju, Chen, Chang, & Wong, 2013; Cheng, Ju, Chen, & Wong, 2012; Ju, Lin, Cheng, Cheng, & Wong, 2013; Ju, Liu, Cheng, & Chang, 2011). Repetitive passive movements decrease spasticity via decreasing stiffness of the mechanical component, especially the muscle–tendon–unit (Cheng et al., 2013). These changes also arouse muscle spindle in detecting its fascicle length (Ju et al., 2013). Consequently, the motor function of the participants improves.

However, research increasingly shows that spasticity and abnormal sensation are not the sole culprits in producing motor dysfunctions such as impaired ambulation in children with CP (Damiano, Martellotta, Sullivan, Granata, & Abel, 2000; McLaughlin et al., 2005). Poor muscle strength is also an important factor that leads to movement dysfunction. Recently, vibratory stimulation was proposed as a new therapeutic modality for the treatment of spasticity and muscle strengthening in participants with upper motor neuron syndrome (Ahlborg, Andersson, & Julin, 2006; Chan et al., 2012; Ness & Field-Fote, 2009). Whole body vibration (WBV), for which the participant stands on a vibrating platform, delivers low-frequency, low-amplitude mechanical stimuli that enter the human body via the feet. The vibrations stimulate the muscle spindles, sending nerve impulses to initiate muscle contractions according to the tonic vibration reflex (Cardinale & Bosco, 2003). Compared to the repetitive passive movement, this WBV protocol adds a muscle strengthening component to the anti-spastic effects.

To date, little research examined the effects of WBV on the lower extremity spasticity and ambulatory function with clinical indicators in children with CP. A recent study examined the effects of a 9-min WBV program on children with CP. The mobility in these children improved, however the only indicator for mobility was the change in self-selected walking speed. No spasticity measurement was performed (Ruck, Chabot, & Rauch, 2010). Another study investigated the vibration effects onto posture and gait (Unger, Jelsma, & Stark, 2013). The vibration used in that study aimed at trunk muscle strengthening. Besides, other than the posture-related measurements, only one gait parameter, the 1-Minute Walk Test, was assessed. The direct effect on spasticity was not evaluated.

Furthermore, one study reported that WBV results in a significant decrease in spasticity in the knee extensors but not in other muscle groups. Participants also showed increased muscle strength in lower legs at a higher isokinetic knee testing speed ( $90^\circ/\text{s}$ ). However, the ambulatory parameters did not change significantly (Ahlborg et al., 2006). Moreover, these findings are from adults with CP. Data from children with CP are relatively scarce. Based on our previous research findings with regard to repetitive passive movement in CP, it is likely that children with CP could benefit from WBV training. Therefore, the aim of this study was to evaluate the effects of WBV on lower extremity spasticity and ambulatory function in children with CP.

## 2. Methods

A complete crossover design was employed in this study. The independent variables were the intervention (WBV or control) and time (pre-WBV = time 1; post-WBV = time 2; 30 min post-WBV = time 3). The effects of WBV were evaluated via variables including active and passive range-of-motion (AROM and PROM) for both knee and ankle joints, relaxation index (RI) assessed by the Wartenburg Pendulum test, Modified Ashworth Scales (MAS) for bilateral knee extensors, timed up-and-go (TUG) and Six Minute Walk Test (6MWT).

### 2.1. Participants

Sixteen children with CP, 9 boys and 7 girls, aged 9.8(2.3) years with diagnoses of either spastic diplegia ( $N = 11$ ) or spastic quadriplegia ( $N = 5$ ) were recruited from a local hospital and a special education school. The inclusion criteria were: add-up MAS scores of bilateral lower extremity greater than two; ability to walk for at least 6 min with or without walking aids; and ability to comprehend commands. Criteria which would affect their muscle tone and ambulatory performance were excluded. Exclusion criteria were: the presence of a progressive neurological, genetic, or metabolic disorder, or a severe concurrent illness or disease not typically associated with CP; lower extremity surgery within the past 6 months; nerve block or botulinum toxin injection within the past 3 months; knee joint range-of-motion limitation greater than  $10^\circ$ ; and epilepsy. One of the authors performed the selection process. Twelve participants walked independently without walking aids and four used 4-wheeled walkers for ambulation. The Institutional Review Board for Human Studies of Chang Gung Memorial Hospital approved this protocol. Written informed consents were obtained from all participants and their legal guardians.

### 2.2. Equipment

The equipment utilized in this study included electrogoniometers (SG110, Biometrics, Ltd., Cwmfelinfach, Gwent, UK) and a WBV platform (AV-001A, Body Green, Taipei, Taiwan). The electrogoniometers, taped on the lateral sides of the knee and ankle joints, were used to measure the degree of flexion and extension. A Biopac MP150 data acquisition unit (Biopac Systems Inc., Santa Barbara, CA, USA) was used for signal registration. The WBV platform could provide vertical vibration with a magnitude of 8–40 Hz and an amplitude of 0.4–2.0 mm.

### 2.3. Protocol

Each subject performed both the WBV and control condition in a counterbalanced order on two separate days, one week apart. The subject was instructed to adhere to the daily routines between the two days. For the WBV condition, participants stood on the vibration platform with the knees slightly flexed at about  $30^\circ$  from full extension. Anterior knee stops and a pelvic belt were used along with the WBV machine for securing their knees and pelvis to provide external support while standing. The participants were instructed to avoid holding on to the supported rail if possible however they were allowed to hold onto the rail if necessary (Ness & Field-Fote, 2009). They were asked to focus on standing with equal weights on both legs. Vibration was delivered at 20 Hz with a vertical displacement of 2 mm. The

WBV was given for 20 min. For the control condition, participants followed the same procedure instead that the vibration machine was not turned on. After the intervention, the participants were moved to the assessment area in wheelchairs to avoid any physical activity on their feet that might influence the vibration effects.

#### 2.4. Assessment

The following measurements were performed before (time 1), immediately after (time 2), and 30 min after (time 3) the intervention, in the order of active and passive range-of-motion measurements (AROM and PROM) for knee/ankle joints, Pendulum test, Modified Ashworth Scale (MAS), timed-up-and-go (TUG), and Six Minute Walk Test (6MWT). The test-retest reliability for each variable was evaluated from two consecutive days before intervention with ten children with spastic cerebral palsy, aged 9.71(1.66) years. Two change scores (Diff2–1/Diff3–1) were calculated from the three time points, with the “before (time 1)” score served as the baseline.

##### 2.4.1. Measurement of joint range-AROM & PROM

The AROM measures the range of movement through which the subject can actively move a joint using the adjacent muscles, whereas the PROM amount of motion at a given joint when the joint is moved by an external force. To measure the flexion and extension, electrogoniometers were secured on the lateral sides of the knee and ankle joints. For the AROM measurement, the subject was first asked to actively move the knee/ankle to full extension and then full flexion. Subject was asked to maintain full extension and flexion for 3 s each and the median angles of the two 3-s intervals were used for AROM calculation. For the PROM condition, their knees/ankles were passively moved by an examiner to the end range for signal registration. The Intraclass correlation coefficients (ICCs) for AROM and PROM were 0.73 and 0.79, respectively.

##### 2.4.2. Measurement of spasticity – Wartenburg Pendulum test

The Pendulum test evaluated muscle tone by using gravity to provoke stretch reflexes of knee extensors during passive swinging of the lower limb. The oscillatory movements of the lower leg were captured by electrogoniometers at the lateral sides of the knee. The relaxation index (RI) was then calculated as follows:  $RI = (\text{starting angle} - \text{first angle}) / (\text{starting angle} - \text{resting angle}) \times 1.6$  (Burridge et al., 2005). The ICC calculated for RI was 0.72. In case of spasticity, the oscillatory amplitudes were significantly less as compared to those with normal muscle tone.

##### 2.4.3. Measurement of spasticity – MAS

The MAS scale is widely used for measuring spasticity. A score of 0–4 was used to assess the resistance of knee extensors to passive movement. The children laid supine with the lower leg hanging freely over the edge and the knee passively extended by the examiner. The examiner moved the subject's knee to a position of maximal flexion over 1 s. A score was then given based on the classification by Bohannon and Smith (1987). The MAS scores of both legs were added up to represent the muscle tone of the subject's lower extremities. The ICC calculated for MAS was 0.92.

##### 2.4.4. Functional measurement – TUG

TUG is a test used to assess a person's mobility. TUG measured the time required for an individual to stand up from a chair with armrests, walk 3 m, turn, walk back to the chair, and sit down. The subject walked with their regular footwear at their preferred speed with or without walking aids. The examiner stayed with the participants to protect their safety at all times. High test-retest reliability (ICC = 0.94) and an adequate validity (Spearman  $r = -.77$ ) for the TUG in children with CP have been reported (Zaino, Marchese, & Westcott, 2004).

##### 2.4.5. Functional measurement – 6MWT

Six-MWT measured the distance walked within 6 min. A 7-m walkway was marked on a level floor. The subject was asked to walk continuously along this walk way, turn around at the end, and continue to cover as much ground as possible over 6 min. The examiner used standardized phrases for speaking

to the patient to avoid improper encouragement and enthusiasm. High test–retest reliability (ICC = 0.98) for the 6MWT in children with CP was revealed (Maher, Williams, & Olds, 2008).

### 2.5. Statistics

SPSS version 17.0 statistical software (SPSS Inc., Chicago, IL, USA) was used to analyze the data. Descriptive statistics were used to calculate participants' demographics. A two-way repeated measures analysis of variance (ANOVA) (condition (2)  $\times$  change score (2)) was performed for all variables. Change scores were used to show the improvement of each dependent variable measured at time 2 and 3 compared to the baseline (time 1). The alpha level was set at 0.05. The relationship between functional improvements and changes in spasticity were examined by correlation analysis.

### 3. Results

Descriptive statistics for the outcome variables in the two conditions on the three measurement times are presented in Table 1. Descriptive and inferential statistics for the change scores are given in Table 2. Significant differences were found in change scores between the treatment and control condition for variables including knee and ankle AROM ( $p = .000$ ), RI ( $p = .000$ ), MAS ( $p = .001$ ), TUG ( $p = .001$ ) and 6MWT ( $p = .000$ ). For MAS ( $p = .007$ ) and 6MWT ( $p = .049$ ) significant differences were also found between the time change scores (2–1) and (3–1). No difference was found in PROM.

The relationships between functional improvements and changes in spasticity were investigated via the correlation of change scores. The results revealed that change in TUG was significantly correlated with change in RI ( $r = -.512$ ,  $p = .042$ ), but not with change in MAS ( $r = -.150$ ,  $p = .580$ ). Change in distance at the Six Minute Walk Test was significantly correlated with change in TUG ( $r = -.700$ ,  $p = .003$ ), but not with change in RI or MAS. No other correlations were significant.

### 4. Discussion

This study evaluated the effect of WBV on lower extremity spasticity and ambulatory function in children with CP. The results revealed that spasticity, measured via the Wartenburg Pendulum test and MAS, decreased. The ambulatory function, tested with TUG and 6MWT, improved significantly after the WBV. The AROM of both knee and ankle joints also increased. Findings of this study demonstrated that the WBV is an effective intervention for controlling spasticity and improving ambulation.

**Table 1**  
Descriptive statistics for outcome variables.

Variables	WBV condition			Control condition		
	Time 1	Time 2	Time 3	Time 1	Time 2	Time 3
KAROM (°)	93.08 (10.29)	98.71 (10.64)	97.41 (10.52)	94.08 (11.03)	93.97 (9.81)	93.37 (8.43)
AAROM (°)	26.98 (6.48)	37.58 (5.30)	37.67 (5.46)	26.98 (6.48)	27.65 (6.43)	27.19 (6.46)
KPROM (°)	104.39 (11.05)	105.32 (10.72)	103.69 (11.23)	104.20 (11.09)	104.21 (10.65)	104.05 (10.53)
APROM (°)	48.82 (7.11)	48.76 (4.69)	49.82 (4.35)	48.82 (7.11)	48.53 (6.91)	48.70 (7.02)
RI	0.59 (0.07)	0.71 (0.07)	0.68 (0.09)	0.59 (0.07)	0.60 (0.08)	0.61 (0.08)
MAS	4.63 (0.96)	3.13 (1.09)	3.56 (1.09)	4.88 (0.96)	4.50 (0.73)	4.75 (0.86)
TUG (s)	13.31 (2.45)	10.45 (1.53)	10.70 (2.47)	13.49 (2.23)	13.09 (1.94)	12.99 (2.03)
6MWT (m)	202.22 (20.63)	219.74 (22.05)	208.35 (24.80)	202.34 (20.59)	200.58 (24.40)	200.69 (20.51)

KAROM/AAROM: knee/ankle active range of motion; KPROM/APROM: knee/ankle passive range of motion; RI: relaxation index; MAS: Modified Ashworth Scale; TUG: timed-up-and-go; 6MWT: Six Minute Walk Test. Values are expressed as mean (SD).

**Table 2**  
Descriptive and inferential statistics for outcome analyses-change scores.

Variables	WBV condition		Control condition		Cohen's $d_{\text{condition}}$		$P_{\text{condition}}$	$P_{\text{change\_score}}$	$P_{\text{interaction}}$
	Change_score <sup>a</sup>		Change_score <sup>a</sup>						
	Diff2-1	Diff3-1	Diff2-1	Diff3-1	Diff2-1	Diff3-1			
KAROM (°)	5.63(3.50)	4.33(2.05)	-0.11(2.53)	-0.71(4.57)	1.880	1.423	0.000**	0.100	0.523
AAROM (°)	10.59(4.40)	10.69(5.31)	0.67(2.45)	0.21(1.86)	2.786	2.634	0.000**	0.659	0.452
KPROM (°)	0.94(2.18)	-0.70(2.93)	0.01(1.56)	-0.15(1.54)	0.491	-0.235	0.750	0.082	0.168
APROM (°)	-0.06(6.01)	-1.00(6.70)	-0.29(1.98)	-0.12(1.90)	0.051	-0.179	0.635	0.181	0.227
RI	0.12(0.65)	0.09(0.06)	0.01(0.05)	0.02(0.04)	0.239	1.373	0.000**	0.217	0.152
MAS	-1.50(1.10)	-1.06(1.12)	-0.38(0.50)	-0.13(0.34)	-1.311	-1.124	0.001**	0.007**	0.270
TUG (s)	-2.86(2.36)	-2.61(1.82)	-0.40(1.29)	-0.51(1.37)	-1.294	-1.304	0.001**	0.826	0.457
6MWT (m)	17.52(15.98)	6.13(7.76)	-1.79(10.29)	-1.65(9.41)	1.437	0.902	0.000**	0.049*	0.025*

Values are expressed as mean(SD). \* $P < .05$ ; \*\* $P < .01$ .

Cohen's  $d$  refers to effect sizes based on the differences between the change scores (2-1 and 3-1) in the WBV and control condition.

<sup>a</sup> Different score: Diff2-1: difference between measurements "time 2" & "time 1"; Diff3-1: difference between measurements "time 3" & "time 1".

These acute effects might promote children's active participation in exercise training and/or therapeutic interventions.

Whole-body vibration was presented to children with CP during supported standing on a vibrating platform. The vibrations are believed to initiate muscle contractions by stimulating muscle spindles and alpha motor neurons. Hence the electromyographic activity increases during WBV, resulting in an effect similar to that of conventional resistance training (Delecluse, Roelants, & Verschueren, 2003). Unlike resistance training which usually needs strong motivation of the participants, vibration therapy might serve as a form of muscle training that is largely independent of the motivation of the patient (Rauch, 2009). WBV provides a great opportunity to patients with neuromuscular disorders who often lack motivation to engage in muscle strengthening exercises.

Previous studies have reported that WBV has anti-spastic effects in patients with neuromuscular diseases (Ahlborg et al., 2006; Chan et al., 2012; Ness & Field-Fote, 2009). Modified Ashworth Scale and the pendulum test are commonly used clinical assessment tools to rate subject's spasticity. The results in this study indicated that after WBV intervention, children with CP demonstrated a decrease in spasticity, as measured with MAS and RI. An electrophysiological study of vibration therapy to local muscles has revealed its effects on spasticity (Liepert & Binder, 2010). The research suggested that vibration effects are mediated through the activation of muscle spindles and transmission by Ia fibers, which enhances cortical excitability of the vibrated muscle. At the same time, the vibration stimuli also reduce activity in antagonistic muscles via reciprocal inhibition and supraspinal inhibition. A more balanced interaction between flexors and extensors can thus be achieved. On the other hand, research on WBV pointed out that it also resulted in a decrease of spasticity (Ahlborg et al., 2006; Chan et al., 2012; Ness & Field-Fote, 2009). With spasticity, the latency of the H-reflex shortens and its amplitude increases. The WBV can facilitate presynaptic inhibition. Presynaptic inhibition of Ia-afferents reduces the release of neurotransmitters to the motoneurons, weakens the effects of Ia-afferents on motoneurons, thus resulting in a decrease of the H-reflex amplitudes. An influence of vibration on mechanical muscle fiber properties may also have contributed to the clinical effect.

The improvement on TUG was significant after WBV intervention and lasted over 30 min. On the other hand, the significant improvement on 6MWT seemed to fade gradually after the cease of the intervention. The correlation analysis revealed that the change in TUG was significantly correlated with the change in RI. The improvement in the latter variable, like the improvement in TUG, was maintained during follow-up period of 30 min. Since spastic muscles exhibit velocity-dependent hyper-tonic reflexes and demonstrate higher passive stiffness, patients with measurable spasticity show reduced angular velocity of the knee during walking and functional performance (Tuzson, Granata, & Abel, 2003). As a consequence, the ambulation function improves as the spasticity decreases. In addition, normalizing muscle tone can ease discomfort, thus improving walking ability and daily functioning. The results on TUG and 6MWT in our study are in line with these considerations. Unlike

TUG, 6MWT might require longer intervention than provided in our study to obtain longer-lasting improvement.

Significant improvement was also noticed in the AROM for the knee and ankle joints, both immediately and 30 min after the intervention. To the authors' knowledge, this was the first study that measured joint ROM change after WBV intervention. The improvement in range of motion can be explained by the stimulation of skin receptors, muscle spindles and the vestibular system (Katusic, Alimovic, & Mejaski-Bosnjak, 2013). In the present study vibratory stimuli were directly delivered to the soles of feet. These mechanical stimuli facilitate the activation of muscle spindles and enhance cortical excitability of the vibrated muscles, leading to joint AROM increase. No such improvements occurred in PROM. PROM is assessed by passively moving a joint to its extreme position. This movement is slow and should not elicit reflex activity. Therefore, the limitation in PROM is mainly due to mechanical constraints, such as the length of musculature surrounding the joint, the flexibility of joint capsule, and the contour of bones, rather than to spasticity.

In addition to its anti-spastic effect, other benefits of WBV are worth mentioning. According to the literature, the oxygen consumption, muscle temperature, skin blood flow and muscle power increase during vibration therapy (Cochrane, Stannard, Sargeant, & Rittweger, 2008; Lohman Iii, Petrofsky, Maloney-Hinds, Betts-Schwab, & Thorpe, 2007). If applied repetitively, positive long-term effects on muscle strength, balance, and bone density have been noted (Rauch, 2009). These results support the clinical application of WBV in CP rehabilitation. Future investigations might focus on finding the optimal combination of frequency and duration of WBV interventions.

This study suggests that a 20-min, 20 Hz WBV intervention can control the spasticity, enhance ambulatory performance and increase AROM in knee and ankle in children with CP. However, a longer follow-up period than the 30 min used in our study should be provided to investigate how long the beneficial effects of a WBV intervention are maintained. Notwithstanding this limitation, however, even on the basis of the short-term improvements demonstrated by us, clinicians might decide to apply WBV prior to exercise or other therapeutic interventions.

Finally, we would like to draw attention to the fact that WBV needs close supervision, in particular when it is offered to patients with physical and/or sensory problems, to safeguard them from losing their balance.

## Conflict of interest

The authors declare that there is no conflict of interest.

## Acknowledgments

The authors would like to thank for the participation of all children, parents, school teachers and staff in this study. This work was supported by the National Science Council (101-2410-H-179-015-), and the Chang Gung Medical Research Program (CMRP), Taiwan (Grant No. CMRPD1D0131).

## References

- Ahlborg, L., Andersson, C., & Julin, P. (2006). Whole-body vibration training compared with resistance training: Effect on spasticity, muscle strength and motor performance in adults with cerebral palsy. *Journal of Rehabilitation Medicine*, 38, 302–308.
- Bohannon, R. W., & Smith, M. B. (1987). Interrater reliability of a modified Ashworth scale of muscle spasticity. *Physical Therapy*, 67, 206–207.
- Burridge, J. H., Wood, D. E., Hermens, H. J., Voerman, G. E., Johnson, G. R., van Wijck, F., et al (2005). Theoretical and methodological considerations in the measurement of spasticity. *Disability and Rehabilitation*, 27, 69–80.
- Cardinale, M., & Bosco, C. (2003). The use of vibration as an exercise intervention. *Exercise and Sport Sciences Reviews*, 31, 3–7.
- Chan, K. S., Liu, C. W., Chen, T. W., Weng, M. C., Huang, M. H., & Chen, C. H. (2012). Effects of a single session of whole body vibration on ankle plantarflexion spasticity and gait performance in patients with chronic stroke: A randomized controlled trial. *Clinical Rehabilitation*, 26, 1087–1095.
- Cheng, H. Y. K., Ju, Y. Y., Chen, C. L., Chang, Y. J., & Wong, A. M. K. (2013). Managing lower extremity muscle tone and function in children with cerebral palsy via eight-week repetitive passive knee movement intervention. *Research in Developmental Disabilities*, 34, 554–561.

- Cheng, H. Y. K., Ju, Y. Y., Chen, C. L., & Wong, M. K. A. (2012). Managing spastic hypertonia in children with cerebral palsy via repetitive passive knee movements. *Journal of Rehabilitation Medicine*, *44*, 235–240.
- Cochrane, D. J., Stannard, S. R., Sargeant, A. J., & Rittweger, J. (2008). The rate of muscle temperature increase during acute whole-body vibration exercise. *European Journal of Applied Physiology*, *103*, 441–448.
- Damiano, D. L., Martellotta, T. L., Sullivan, D. J., Granata, K. P., & Abel, M. F. (2000). Muscle force production and functional performance in spastic cerebral palsy: Relationship of cocontraction. *Archives of Physical Medicine and Rehabilitation*, *81*, 895–900.
- Delecluse, C., Roelants, M., & Verschuere, S. (2003). Strength increase after whole-body vibration compared with resistance training. *Medicine and Science in Sports and Exercise*, *35*, 1033–1041.
- Ju, Y. Y., Lin, J. K., Cheng, H. Y. K., Cheng, C. H., & Wong, A. M. K. (2013). Rapid repetitive passive movement promotes knee proprioception in the elderly. *European Review of Aging and Physical Activity*, *10*, 133–139.
- Ju, Y. Y., Liu, Y. C., Cheng, H. Y. K., & Chang, Y. J. (2011). Rapid repetitive passive movement improves knee proprioception. *Clinical Biomechanics*, *26*, 188–193.
- Katusic, A., Alimovic, S., & Mejaski-Bosnjak, V. (2013). The effect of vibration therapy on spasticity and motor function in children with cerebral palsy: A randomized controlled trial. *NeuroRehabilitation*, *32*, 1–8.
- Koman, L. A., Smith, B. P., & Shilt, J. S. (2004). Cerebral palsy. *Lancet*, *363*, 1619–1631.
- Krigger, K. W. (2006). Cerebral palsy: An overview. *American Family Physician*, *73*, 91–100.
- Liepert, J., & Binder, C. (2010). Vibration induced effects in stroke patients with spastic hemiparesis – A pilot study. *Restorative Neurology and Neuroscience*, *28*, 729–735.
- Lohman III, E. B., Petrofsky, J. S., Maloney-Hinds, C., Betts-Schwab, H., & Thorpe, D. (2007). The effect of whole body vibration on lower extremity skin blood flow in normal subjects. *Medical Science Monitor*, *13*, CR71–CR76.
- Maher, C. A., Williams, M. T., & Olds, T. S. (2008). The six-minute walk test for children with cerebral palsy. *International Journal of Rehabilitation Research*, *31*, 185–188.
- McLaughlin, J. F., Felix, S. D., Nowbar, S., Ferrel, A., Bjornson, K., & Hays, R. M. (2005). Lower extremity sensory function in children with cerebral palsy. *Pediatric Rehabilitation*, *8*, 45–52.
- Ness, L. L., & Field-Fote, E. C. (2009). Effect of whole-body vibration on quadriceps spasticity in individuals with spastic hypertonia due to spinal cord injury. *Restorative Neurology and Neuroscience*, *27*, 621–631.
- Rauch, F. (2009). Vibration therapy. *Developmental Medicine and Child Neurology*, *51*, 166–168.
- Ruck, J., Chabot, G., & Rauch, F. (2010). Vibration treatment in cerebral palsy: A randomized controlled pilot study. *Journal of Musculoskeletal Neuronal Interactions*, *10*, 77–83.
- Tuzson, A. E., Granata, K. P., & Abel, M. F. (2003). Spastic velocity threshold constrains functional performance in cerebral palsy. *Archives of Physical Medicine and Rehabilitation*, *84*, 1363–1368.
- Unger, M., Jelsma, J., & Stark, C. (2013). Effect of a trunk-targeted intervention using vibration on posture and gait in children with spastic type cerebral palsy: A randomized control trial. *Developmental Neurorehabilitation*, *16*, 79–88.
- Zaino, C. A., Marchese, V. G., & Westcott, S. L. (2004). Timed up and down stairs test: Preliminary reliability and validity of a new measure of functional mobility. *Pediatric Physical Therapy*, *16*, 90–98.